



**DSC Lab**

**BETA-STIM<sup>TM</sup>**

**Label**

## WARNINGS:

- Keep in a cool, dry, and dark place.
- Do not use product after expiry date.
- Do not store near pesticides.

Although this product has been tested under a variety of conditions, DSC Lab CC does not warrant that it will be efficacious under all conditions, as the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. DSC Lab CC furthermore does not accept any responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow the label instructions. Consult DSC Lab CC in case of any uncertainty.

## PRECAUTIONS:

- P264 – Wash hands thoroughly after handling.
- P270 – Do not eat. Drink or smoke while using this product.
- P405 – Store locked up.

## FIRST AID:

First-aid measures after inhalation: If person is unconscious seek emergency medical attention. If person is conscious remove to fresh air and call a physician.

First-aid measures after skin contact: Should pose no danger. Wipe or rinse off using water.

First-aid measures after eye contact: Flush eyes with copious amounts of water. Call physician if needed.

First-aid measures after ingestion: Ingestion, though not advised, should pose no danger. Drink water. Seek medical attention if any adverse effects are noted.

## USE RESTRICTIONS:

The product should not be used for any other purpose or in any other manner contrary to the information supplied on the product label.

## DIRECTIONS FOR USE:

Use only as directed.

- Shake well before use.
- Do not use the product after the expiry date.
- Use product within 12 hours after opening the container.
- Do not leave the product in the sun when not using. These are living organisms that are sensitive to sunlight.

BETA-STIM is registered for use on row crops (maize, grains and related crops), soybean and other legume crops, as well as oilseed crops (sunflower).

APPLICATION METHOD	DOSAGE	RECOMMENDATIONS
Seed Treatment	200 ml BETA-STIM p/50 kg seed	Apply directly to seed. Can be diluted with water. Water volume is determined by product and seed type.
In-furrow Application	800 ml BETA-STIM p/ha in-furrow	Pre-mix product with 5 ℓ water before adding to application tank. Ensure thorough agitation. Apply in 25 – 250 ℓ of water per hectare.
Soil and Irrigation Application	400 – 800 ml BETA-STIM p/ha	Pre-mix with 5 ℓ water. Mix thoroughly before applying through irrigation system.