



DSC Lab
TRICHODERMATM
Label

WARNINGS:

- Keep in a cool, dry, and dark place.
- Do not use product after expiry date.
- Do not store near pesticides.

Although this product has been tested under a variety of conditions, DSC Lab CC does not warrant that it will be efficacious under all conditions, as the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. DSC Lab CC furthermore does not accept any responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow the label instructions. Consult DSC Lab CC in case of any uncertainty.

PRECAUTIONS:

- P264 – Wash hands thoroughly after handling.
- P270 – Do not eat. Drink or smoke while using this product.
- P405 – Store locked up.

FIRST AID:

First-aid measures after inhalation: If person is unconscious seek emergency medical attention. If person is conscious remove to fresh air and call a physician.

First-aid measures after skin contact: Should pose no danger. Wipe or rinse off using water.

First-aid measures after eye contact: Flush eyes with copious amounts of water. Call physician if needed.

First-aid measures after ingestion: Ingestion, though not advised, should pose no danger. Drink water. Seek medical attention if any adverse effects are noted.

USE RESTRICTIONS:

The product should not be used for any other purpose or in any other manner contrary to the information supplied on the product label.

DIRECTIONS FOR USE:

Use only as directed.

- Shake well before use.
- Do not use the product after the expiry date.
- Use product within 12 hours after opening the container.
- Do not leave the product in the sun when not using. These are living organisms that are sensitive to sunlight.

TRICHODERMA is registered for use on row crops (maize, grains, sunflower and related crops) as well as soybean and other legume crops.

APPLICATION METHOD	DOSAGE	RECOMMENDATIONS
Seed Treatment	200 g TRICHODERMA p/50 kg seed	Apply directly to seed. Suspend product in as much water as is necessary to form slurry. Mix the slurry with seed until all seeds are evenly coated. Let dry and plant seed as soon as seed is manageable.
In-furrow Application	800 g TRICHODERMA p/ha in-furrow	Apply as a suspension on the seed in the planter furrow. Use as much water as is convenient. Ensure thorough agitation in the application tank.
Soil and Irrigation Application (drip or pivot irrigation)	800 g TRICHODERMA p/ha	Mix thoroughly before applying through irrigation system. Use as much water as is necessary.