

Chapter 8

The Presence of Endometriosis in the Human Fetus



Pietro G. Signorile and Alfonso Baldi

Introduction

Endometriosis is characterized by the growth of endometrial glands and stroma at extrauterine sites, most frequently over visceral and peritoneal surfaces within the female pelvis [1]. It is a very common gynecological disorder present in up to 10% of women of reproductive age [2]. The incidence rises to 30% in patients with difficulties in conceiving [3]. Deep infiltrating endometriosis is a subset of endometriosis where the lesions are predominantly located under the peritoneal surface and is associated with intense pelvic pain symptoms [4]. Endometriosis is generally accompanied by chronic pelvic pain, adhesion formation, and infertility. It has been calculated that endometriosis is responsible for more than 100,000 hysterectomies each year in the United States with significant annual health-care costs attributable to this disease [3]. Moreover, the symptoms of this disease are mostly nonspecific and are very similar to those associated with other chronic pain disorders. As a result, in a great majority of cases, the definitive diagnosis is reached after several years and only by invasive surgical procedures, often with an incredible time interval between the onset of the symptoms and final diagnosis of 8–12 years [2, 3]. Original noninvasive approaches for the diagnosis of endometriosis have been recently proposed by our research group; however, this therapy is not yet standard of care [5–8]. Endometriosis remains a significantly underdiagnosed and under-treated disease and is considered a “social disease” since it has an important socio-economic impact in view of the costs for the diagnosis and treatment, the loss of economic performance of the patients, and the negative impact on quality of life and capability of conceiving [2, 3].

P. G. Signorile (✉) · A. Baldi
Fondazione Italiana Endometriosi, Rome, Italy
e-mail: research@endometriosi.it

Pathogenesis of Endometriosis

Thanks to the work of Benagiano and Brosen and that of Knapp, the steps in the discovery and characterization of endometriosis in the history of medicine are now very well defined [9, 10]. Endometriosis has been a known entity for more than two centuries. In light of the significant social and economic impact this disease is known to have, it seems impossible that the pathogenesis of endometriosis has not been definitively elucidated. Currently, several pathogenetic theories have been proposed to explain the development and establishment of endometriosis. Table 8.1 summarizes the most commonly accepted theories for the histogenesis of endometriosis.

The most easily understood and widely accepted theory for the histogenesis of endometriosis is that of retrograde menstruation/transplantation, proposed one century ago by Sampson [2]. This theory hypothesizes that at menstruation, some effluent flows retrograde through the lumen of the Fallopian tubes into the peritoneal cavity, causing the adhesion and growth of endometrial fragments. This mechanism that considers endometriosis simply an autotransplant of normal endometrial tissue in an ectopic location in the organism explains the most common sites of endometriosis [2, 3]. Moreover, retrograde menstruation is a common phenomenon with more than 90% of women having blood in their pelvis at the time of menstruation [1]. Despite the fact that researchers have tried for decades to confirm this mechanism of histogenesis for endometriosis, no conclusive evidence has been produced in favor of this theory. More importantly, it fails to explicate the presence of endometriosis in such remote areas outside the peritoneal cavity, as the lungs, skin, lymph nodes, and breasts [2, 3]. Moreover, it is not an acceptable pathogenetic mechanism for endometriosis described in early puberty and in newborns [11], as well as in women affected by Mayer-Rokitansky-Küster-Hauser, a syndrome characterized by congenital aplasia of the uterus and the upper part of the vagina [12]. Finally, it cannot be considered a valid mechanism in the event of endometriosis in male, which is a rare but very well-described phenomenon with a total of 17 cases reported in the literature to date [13]. Nevertheless, elegant observations by Redwine propose that endometriotic tissue lacks characteristics of an autotransplant [14].

Table 8.1 Different theories proposed for the etiology of endometriosis

Theory	Mechanism proposed
Retrograde menstruation/transplantation	Retrograde menstruation allows implantation of endometrial glands into the peritoneal cavity.
Coelomic metaplasia	Endometriosis arises in the pelvis or elsewhere by endometrial metaplasia of peritoneal mesothelium or other cell types.
Lymphatic and vascular dissemination	Spread of endometrial cells happens by lymphatic or hematogenous vessels.
Circulating stem cells	Transient pluripotent hematopoietic stem cells could differentiate into endometriotic tissue at different anatomical sites.
Embryonic cell remnants	Endometriosis originates from embryonic rests of the Mullerian ducts and Wolffian ducts.

The coelomic metaplastic theory suggests that endometriosis in the pelvis and elsewhere is caused by endometrial metaplasia of the peritoneal serosa or serosa-like structures, perhaps induced by environmental factors [15]. This theory would explain the cases where retrograde menstrual flow is impossible.

The lymphatic and vascular dissemination theory suggests the spread of endometrial cells by lymphatic or hematogenous vessels [15].

The theory of circulating stem cells claims that transient pluripotent hematopoietic stem cells could differentiate into endometriotic tissue at different anatomical sites [15].

The theory of embryonic cell remnants postulates that endometriosis originates from embryonic rests of the Mullerian ducts and Wolffian ducts. The Müllerian ducts, indeed, give rise to the female reproductive tract. This organogenesis is controlled by complex spatiotemporal molecular pathways, including the anti-Müllerian hormone signaling [15]. Aberrant differentiation or migration of the Müllerian ducts during embryogenesis could spread cells in their migratory pathway across the posterior pelvic floor, thus clarifying the observation that endometriosis is commonly found in the cul-de-sac, uterosacral ligaments, and medial broad ligaments.

This mechanism of histogenesis was proposed by pioneer scientists of this disease in the late nineteenth and twentieth century but inexplicably forgotten after the advent of Sampson’s retrograde menstruation theory [9, 10]. Table 8.2 summarizes the most important observations supporting the embryological theory in chronological order as clearly described by the works of Benagiano and Brosens and Knapp [9, 10]. Further supporting evidence of the fetal origin is the observation that the cells of pubertal or postpubertal clinical endometriosis maintain some typical characteristics of fetal endometrium cells, such as ontogenic resistance to progesterone [16]. This characteristic is, obviously, not present in the mature adult endometrium.

Recently, work from our research group and others have demonstrated the presence of ectopic endometrium in a significant number of human female fetuses [17–19]. These scientific evidences clearly support the embryogenetic theory. In this

Table 8.2 Observations supporting the embryological origin of endometriosis, performed by pioneer scientists of this disease in the late nineteenth and twentieth century, as described in the work of Benagiano and Brosens [10] and Knapp [9]

Author	Mechanism proposed	References
Von Reckinglausen 1893	Wolffian origin	Dtsch med Wochenschr 1893; 46: 825.
Orloff 1895	Embryonic cells	Zeitschr Heilkunde 1895; 5: 121.
Pick 1897	Mesonephric origin	Arkiv f Gynäk 1897; 54: 119.
Kossman 1897	Mullerian origin	Archiv f Gynäk 1897; 54: 359.
Mayer 1903	Epithelial heterotopy	Z Geburtshilfe Gynäkol 1903; 49: 32.
Schikele 1904	Mesonephric origin	Zentralbl Allg Pathol Anat 1904; 15: 261
Cullen 1908	Mucosal theory	WB Saunders 1908
Frankl 1911	Mullerian origin	Arkiv f Gynäkol 1911; 93: 659
Lockyer 1918	Mullerian origin	MacMillan and Co, 1918

chapter, we describe the most salient data produced by our research group and other scientists regarding fetal endometriosis in order to better determine the real biological impact of this phenomenon.

Fetal Endometriosis

Our research group has demonstrated the presence of ectopic endometrium in a significant number of human female fetuses (10 in 101 cases) analyzed by autopsy in three different works [17–19]. These structures were found outside the uterine cavity and could not be attributed to any normal anatomical formation. In particular, the anatomical sites of these endometrial structures were in the mesenchymal tissue close to the posterior wall of the uterus, in the proximity of the Douglas pouch, in the rectovaginal septum, in the rectal tube at the level of muscularis propria, and in the wall of the uterus. Interestingly, all of these anatomical sites are very well-known locations for endometriosis in women [1]. Immunohistochemical studies were utilized to assist in the evaluation and characterization of these endometriotic structures. These organoid lesions demonstrated positive staining for CA-125, cytokeratin 7, and estrogen receptor in the epithelial component, while the stromal cells displayed positive staining for both CD-10 and estrogen receptor. Fetal endometrium was also evaluated in these patients and revealed identical staining patterns. The exact anatomical distributions of all the endometriosis-like structures found in these three works are depicted in detail in Fig. 8.1. An example of the histological and immunohistochemical appearance of this ectopic endometrium is reported in Figs. 8.2, 8.3, 8.4, and 8.5. Based on the anatomical location, and on the histological and immunohistochemical characteristics, these structures must be ascribed to endometrial tissue, dislocated outside the uterine cavity during the earlier steps of organogenesis and displaying a molecular immunophenotype identical to that of the endometrium present in the uterus. To the best of our knowledge, these observations have been the first direct and systematic demonstration of the theory of embryonic cell remnants as the cause of endometriosis.


After our data was published, subsequent studies also confirmed the presence of endometriotic structure in female fetuses. In the work by de Jolinière et al., the reproductive organs of seven female fetuses were analyzed at autopsy [20]. In two out of seven fetuses, ectopic endometrial glands were found in the myometrium, while several ectopic endometrial glands surrounded by stroma were found in the uterine broad and ovarian ligaments and under the fallopian tube serosa in six fetuses. These glandular structures expressed positive staining for estrogen and progesterone receptors, while the stromal components displayed positive staining for CD-10 and vimentin [20].

Nevertheless, Schuster and Mackeen have reported a case of fetal endometriosis diagnosed as a large fetal pelvic mass at 35 weeks of gestation [21]. The mass was surgically removed on the second day of life, and histological examination of the specimen confirmed the diagnosis of cystic endometriosis of the left ovary [21].

Fig. 8.1 Anatomical distribution of the ectopic endometrium found in the female human fetuses in the works by Signorile et al. (see references 24–26)

Representation of the pelvic organs of a female fetus at around 25 weeks of gestation, displaying the anatomical location of the endometriotic structures. The different locations are indicated by asterisks in the proximity of the Douglas pouch, in the mesenchymal tissue close to the posterior wall of the uterus, in the rectal tube at the level of muscularis propria, in the wall of the uterus, and in the rectovaginal septum

Abbreviations used: an anus, co coccyx, va vagina, re rectum, sc spinal column, ut uterus, bl bladder (Modified from Signorile et al. [18])



If we analyze the scientific literature preceding the above-described work on fetal endometriosis, we find some anecdotal scientific observations [22, 23]. Moreover, the presence of endometriosis in the fetus was hypothesized, but not demonstrated, by Batt et al. with the mullerianosis theory, even if this phenomenon was considered as different from endometriosis [24]. Based on our data and the observation of others regarding the presence of ectopic endometrium in the fetus, we believe these endometriotic structures remain quiescent and asymptomatic until puberty, at which time the hormonal inputs result in activation and, consequently, the onset of the symptoms of endometriosis [15]. The molecular mechanisms responsible for the histogenesis of these ectopic endometrial structures, as well as their survival until puberty, are largely unknown. As far as pathogenesis is concerned, it is possible to hypothesize that complex molecular mechanisms, also regulated according to a precise spatiotemporal scheme, can be altered by abnormal inputs that act on an adequate genetic background in a specific organogenesis window of time. We consider this phenomenon as an alteration of the fine-tuning of female genital structures organogenesis caused by genetic and epigenetic factors that would cause disruption of some organizational events associated with




Fig. 8.2 Histological and immunohistochemical appearance of ectopic endometrium in a female human fetus of 25 weeks

The picture shows an endometrial structure in the rectovaginal septum; in the inset, the immunohistochemical expression of estrogen of this structure at higher magnification is depicted. (Modified from Signorile et al. [17])



Fig. 8.3 Histological and immunohistochemical appearance of ectopic endometrium in a female human fetus of 24 weeks

The image shows an endometrial structure in the proximity of the Douglas pouch; in the inset, the immunohistochemical expression of estrogen receptor of this structure at higher magnification is depicted. (Modified from Signorile et al. [17])




Fig. 8.4 Histological and immunohistochemical appearance of ectopic endometrium in a female human fetus of 18 weeks

The picture shows an endometrial structure in the rectal tube at the level of muscularis propria; in the inset, the immunohistochemical expression of CA-125 of this structure at higher magnification is depicted. Note that the epithelium of the rectum is negative for CA-125. (Modified from Signorile et al. [17])




Fig. 8.5 Histological and immunohistochemical appearance of ectopic endometrium in a female human fetus of 16 weeks

The image shows an endometrial structure in the mesenchymal tissue close to the posterior wall of the uterus; in the inset, the immunohistochemical expression of CA-125 of this structure at higher magnification is depicted. Note that in the wall of the primitive myometrium is present a little group of endometrial cells positive for CA-125 (indicated by an asterisk) that could represent a primitive nest of adenomyosis. (Modified from Signorile et al. [17])

development of the normal neonatal uterine wall [25]. The observation that endometriosis is significantly higher in patients affected by uterine malformations, such as Müllerian anomalies, anogenital distance, myometrial structural alterations, and other genital anomalies, is indirect support of this proposed mechanism [25]. Recently, works by Makiyan [26] have suggested that the origin of endometriosis is from primordial germ cells; this observation further supports the presence of endometriosis in the fetus.

If we consider the fact that the most important hormone in the female genital tract morphogenesis is estrogen, it is likely that altered estrogenic input may be one of the factors responsible for the histogenesis of fetal endometriosis [25]. There are important epidemiological and experimental studies that link the onset of endometriosis, as well as other changes in the female genital system, to exposure in utero to endocrine disruptors, substances capable of mimicking the action of the hormone estrogen. In this regard, it is important to remember the epidemiological work of Missmer, which demonstrated a significantly higher number of cases of endometriosis in female patients exposed in utero to diethylstilbestrol [27] as well as experimental data from our group showing an endometriosis-like phenotype in mice exposed in utero to the endocrine disruptor bisphenol [28]. Finally, through experimental approaches of genomics, we have been able to reveal a specific pattern of gene expression in the tissue of endometriotic structures compared to normal endometrial tissue. This specific gene expression pattern essentially concerned genes involved in embryogenesis and was not modified by the phases of the hormone cycle [15].

Conclusions

Endometriosis is a multifactorial disease with many-sided features, and accordingly, the pathogenesis could be slightly different in different cases. The proposed theories of pathogenesis are perhaps not mutually exclusive and may be interrelated. Nevertheless, the recent findings reviewed in this chapter lend a compelling argument to the embryogenesis theory especially when compared to the retrograde menstruation theory. The clinical and therapeutic implications are relevant. In particular, recurrence of the disease should not be ascribed to retrograde menstruation but rather to an incomplete surgery due to the presence of microscopic foci and/or at a different timing in the growth of the various foci in the same patient. This last phenomenon is very common in diseases induced by endocrine disruptors and is probably due to individual susceptibility. Even more important, treatment of the disease with synthetic estrogens or selective estrogen receptor modulators chemical compound is, indeed, effective in reducing the symptoms but could cause the growth of microscopic lesions and exacerbate the disease. A more complete understanding of the molecular mechanisms responsible of the pathogenesis of endometriosis could, hopefully, individuate suitable therapeutic targets for this still “uncurable” disease.

Acknowledgments This work was supported by a grant from Fondazione Italiana Endometriosis.

References

1. Baldi A, Campioni M, Signorile PG. Endometriosis: pathogenesis, diagnosis, therapy and association with cancer. *Oncol Rep.* 2008;19:843–6.
2. Giudice LC, Kao LC. Endometriosis. *Lancet.* 2004;364:1789–99.
3. Bulun SE. Endometriosis. *N Engl J Med.* 2009;360:268–79.
4. Signorile PG, Campioni M, Vincenzi B, D'Avino A, Baldi A. Rectovaginal septum endometriosis: an immunohistochemical analysis of 62 cases. *In Vivo.* 2009;23:459–64.
5. Signorile PG, Baldi A. Serum biomarker for the diagnosis of endometriosis. *J Cell Physiol.* 2014;229:1731–5.
6. Signorile PG, Baldi A. Supporting evidences for potential biomarkers of endometriosis detected in peripheral blood. *Data Brief.* 2015;5:971–4.
7. Signorile PG, Baldi A. Prototype of multiplex bead assay for quantification of three serum biomarkers for in vitro diagnosis of endometriosis. *J Cell Physiol.* 2016;231:2622–7.
8. Signorile PG, Baldi A. A tissue specific magnetic resonance contrast agent, Gd-AMH, for diagnosis of stromal endometriosis lesions: a phase I study. *J Cell Physiol.* 2015;230:1270–5.
9. Knapp VJ. How old is endometriosis? Late 17th and 18th century European descriptions of the disease. *Fertil Steril.* 1999;72:10–4.
10. Benagiano G, Brosens I. History of adenomyosis. *Best Pract Res Clin Obstet Gynaecol.* 2006;20:449–63.
11. Marsh EE, Laufer MR. Endometriosis in premenarcheal girls who do not have an associated obstructive anomaly. *Fertil Steril.* 2005;83:758–60.
12. Balci O, Karatayli R, Capar M. An incidental coexistence of Mayer-Rokitansky-Kuster syndrome with pelvic ectopic kidney and perirenal endometrioma. *Saudi Med J.* 2008;29:13450–1341.
13. Rei C, Williams T, Feloney M. Endometriosis in a man: a rare source of abdominal pain: a case report and review of the literature. *Case Rep Obstet Gynecol.* 2018;2018:2083121.
14. Redwine DB. Was Sampson wrong? *Fertil Steril.* 2002;78:686–93.
15. Signorile PG, Baldi A. Endometriosis: new concepts in the pathogenesis. *Int J Biochem Cell Biol.* 2010;42:778–80.
16. Brosens I, Muter J, Gargett CE, Puttemans P, Benagiano G, Brosens JJ. The impact of uterine immaturity on obstetrical syndromes during adolescence. *Am J Obstet Gynecol.* 2017;217:546–55.
17. Signorile PG, Baldi F, Bussani R, D'Armiento MR, De Falco M, Baldi A. Ectopic endometrium in human fetuses is a common event and sustains the theory of müllerianosis in the pathogenesis of endometriosis, a disease that predisposes to cancer. *J Exp Clin Cancer Res.* 2009;28:49.
18. Signorile PG, Baldi F, Bussani R, D'Armiento MR, De Falco M, Boccellino M, Quagliuolo L, Baldi A. New evidence sustaining the presence of endometriosis in the human foetus. *Reprod Biomed Online.* 2010;21:142–7.
19. Signorile PG, Baldi F, Bussani R, Viceconte R, Bulzomi P, D'Armiento M, D'Avino A, Baldi A. Embryologic origin of endometriosis: analysis of 101 human female fetuses. *J Cell Physiol.* 2012;227(4):1653–6.
20. Bouquet de Jolinière J, Ayoubi JM, Lesec G, Validire P, Goguvin A, Gianaroli L, Dubuisson JB, Feki A, Gogusev J. Identification of displaced endometrial glands and embryonic duct remnants in female fetal reproductive tract: possible pathogenetic role in endometriotic and pelvic neoplastic processes. *Front Physiol.* 2012;3:444.
21. Schuster M, Mackeen DA. Fetal endometriosis: a case report. *Fertil Steril.* 2015;103:160–2.
22. Fujii S. Secondary Müllerian system and endometriosis. *Am J Obstet Gynecol.* 1991;165:218–25.
23. Redwine DB. Mülleriosis: the single best fit model of origin of endometriosis. *J Reprod Med.* 1998;33:915–20.
24. Batt RE, Smith RA, Buck Louis GM. Müllerianosis. *Histol Histopathol.* 2007;22:1161–6.

25. Signorile PG, Baldi A. New evidence in endometriosis. *Int J Biochem Cell Biol.* 2015;60:19–22.
26. Makiyan Z. Endometriosis origin from primordial germ cells. *Organogenesis.* 2017;13:95–102.
27. Missmer SA, Hankinson SE, Spiegelman D, Barbieri RL, Michels KB, Hunter DJ. In utero exposure and the incidence of endometriosis. *Fertil Steril.* 2004;82:1501–8.
28. Signorile PG, Spugnini EP, Mita L, Mellone P, D'Avino A, Bianco M, Diano N, Caputo L, Rea F, Viceconte R, Portaccio M, Viggiano E, Citro G, Pierantoni R, Sica V, Vincenzi B, Mita DG, Baldi F, Baldi A. Pre-natal exposure of mice to bisphenol A elicits an endometriosis-like phenotype in female offspring. *Gen Comp Endocrinol.* 2010;168:318–25.